

SEARCH FOR TeV COUNTERPARTS IN GAMMA RAY BURSTS

P.J.Boyle¹, J.H.Buckley², A.M. Burdett³, J.Bussóns Gordo¹, D.A.Carter-Lewis⁴, M.Catanese⁴, M.F.Cawley⁵, V.Connaughton⁶, D.J.Fegan¹, J.P.Finley⁷, J.A.Gaidos⁷, K.Harris², A.M.Hillas³, R.C.Lamb⁸, F.Krennrich⁴, R.W.Lessard⁷, C.Masterson¹, J.E.McEnery¹, G.Mohanty⁴, N.A.Porter¹, J.Quinn^{1,2}, A.J.Rodgers³, H.J.Rose³, F.W.Samuelson⁴, G.H.Sembroski⁷, R.Srinivasan⁷, T.C.Weekes², J.Zweerink⁴,

¹ *Physics Department, University College, Dublin, Ireland*

² *Whipple Observatory, Harvard-Smithsonian CfA, USA*

³ *Physics Department, University of Leeds, United Kingdom*

⁴ *Department of Physics and Astronomy, Iowa State University, USA*

⁵ *Physics Department, St.Patrick's College, Maynooth, Ireland*

⁶ *Marshall Space Flight Center, Alabama, USA*

⁷ *Department of Physics, Purdue University, Indiana, USA*

⁸ *Space Radiation Laboratory, Caltech, Pasadena, USA*

ABSTRACT

Based on BACODINE network notification the Whipple Observatory gamma-ray telescope has been used to search for the delayed TeV counterpart to BATSE-detected gamma-ray bursts. In the fast slew mode, any point in the sky can be reached within two minutes of the burst notification. The search strategy, necessary because of the uncertainty in burst position and limited FOV of the camera, is described.

INTRODUCTION

The Whipple collaboration has an active observational program which is dedicated to searching for TeV counterparts of classical gamma ray bursts. Since May 1994, the Whipple Observatory 10 meter gamma-ray telescope (Cawley, et al., 1990) has made 16 follow-up observations of BATSE burst notifications, but to date no positive identifications have been made (Connaughton, et al., 1995, Connaughton, et al., 1997a). The field of view of our current instrument is limited to 3.5 degrees and the error box for the BATSE position of gamma-ray bursts is 10 degrees in diameter. Therefore, more than a single pointing has to be used for the TeV follow up after a gamma-ray burst. The current observation technique is to take a 28 minute exposure at the initial location and 4 more exposures 3 degrees away, with a final exposure taken again at the initial position as in Figure 1a (Connaughton, et al., 1997a). In order to find any associated TeV component a two dimensional (2D) search strategy has been developed in order to locate the precise region of emission (Akerlof, et al., 1993, Connaughton, et al., 1997a, Lessard, et al., 1997).

Assuming gamma ray bursts have an associated TeV component, it is uncertain as to just how long after the BATSE notification of a gamma-ray burst that this emission would last. At present, we cover less than 50% of the actual BATSE error box in 2.5 hours and of the 16 follow-ups observed to date, only 6 had full observations of all positions (Table 1). All other follow-ups were terminated after an hour due to sun/moonrise or the position being too low to track. Owing to the uncertainty in the timescale of bursts and the lack of coverage of the whole error box, a non-detection cannot be used as an argument against the possibility of gamma-ray burst emission extending up to TeV-emission energies.

Date	BATSE Trig. ^a	BACODINE Intensity Cts/s ^b	Difference BACO- Hunts. (deg) ^c	Durat. BATSE (s) ^d	Whipple delay (min) ^e	Elev- ation (deg) ^f	Cycles observed (1-6) ^g
940516	2980	630	N/A	N/A	19	24-14	1 (1 hour)
950208	3408	778	1.64	N/A	16	N/A	1,2
950405	3494	704	9.5	N/A	8	27-33	1,2,3+cont
950524	3598	8726	1.37	6	5	56-31	complete
950625	3649	1661	3.84	40	18	28-41	1-6
950701	3658	9134	2.3	15	56	41-69	complete
951117	3909	1955	10.1	25	5	30-24	1 (1 hour)
951119	3911	801	6.98	60	20	45-54	complete
951124	3918	1231	5.98	150	2	59-76	complete
951220	4048	6698	2.88	17	27	71-37	complete
960521	5467	2919	9.13		14	80-68	1,2
961017	5634	3193	2.74	41	7	28-23	complete
961111 ^h	5665	496	-	-	4	54-80	1-4
961206	5706	1312	6.4	1.4	5	26-22	1-3
970304	6113	8842	3.1	11	3	28-42	1-4
970308	6117	1120	13.6	2	4	43-72	complete

^aTrigger number of BATSE burst

^bNumber of counts/s measured by BATSE in the first 1 or 2 seconds of the burst.

^cAngular difference between the burst coordinates provided through BACODINE and the final estimated burst position from Huntsville.

^dDuration of the burst seen by BATSE.

^eLength of time that elapsed between sending the BACODINE message and the start of the first Whipple run.

^fAverage elevations of the first and last positions covered by the 10-metre telescope.

^gWhipple coverage of burst area. The numbers listed indicate the positions covered in the observations. The 6 positions are described in the text and are shown in Figure 1a. An entry of "complete" means all 6 positions and their control observations were completed.

^hTrigger #5665 was a terrestrial trigger

RASTER SEARCH TECHNIQUE

In order to achieve a better coverage of the 10 degree error box we now scan a circular region around the BATSE burst position with a constant angular velocity. A raster scan across parallel lines in declination (Figure 1b) with a spacing of 1 degree results in complete coverage of the BATSE error box. A continuous motor speed of 0.1 degree/s facilitates a scan of the whole error box within 400 seconds. The fast scanning method has the advantage that we may give a flux upper limit within the first 300-400 seconds or less depending on the final improved BATSE burst position. Although the sensitivity within the first 300 seconds is not great, a burst of the calibre of 910503 or 930131 would be easily detected if the spectrum extends from the GeV region up to TeV energies (Hurley 1996). The time scale of those two bursts was 84 and 100 seconds respectively and only if the burst positions are close to the actual telescope position can a resonable coverage of the BATSE error box be achieved.

Recent improvements have lowered the time for response to a BACODINE notification of the Whipple telescope to within 15 seconds of the BATSE trigger. A new tracking system for the telescope will be




Fig. 1: (a): Current Response to BACODINE notifications. Each circle represents the field of view of the 10-meter reflector at the 5 different positions following a BATSE burst. (b): Raster Scan response. The reflector is moved at a constant velocity continuously along the track to cover the BATSE error box.

implemented in the fall of 1997, making the most distant point on the sky accessible within 2 minutes. A burst seen by BATSE and EGRET on 940217 lasted for some 1.5 hours at GeV energies. This class of gamma-ray burst might easily have been missed by the previous method but it could now be observed with a reasonable sensitivity (9 minutes effective observation time assuming a 3 degree FOV and a 10 degree error box).

The result of each raster scan will be analysed and displayed by a quicklook analysis to search for a strong excess. If there is no significant excess this scanning process can be continued for 2-5 hours if possible. If something significant is apparent, the telescope will be pointed at this position.

ANALYSIS

The Whipple high energy gamma-ray telescope images Čerenkov light from air showers onto an array of 151 photomultipliers covering a 3.5 degree field of view. The imaging technique uses information of the angular spread and orientation to reject more than 99.7% of the cosmic-ray induced showers while retaining over 50% of the possible gamma-ray induced events. For discrete objects the source is placed at the center of the field of view and candidate gamma-ray events (CGRE) are selected on the basis of the a combination of (a) image shape (*length & width*) and (b) orientation of the major axis of the shower (Reynolds, et al., 1993). However, as a burst source has an unknown position a 2-D search can determine the incident direction of the CGRE. The search makes use of the orientation, elongation and asymmetry of the image. Monte Carlo studies have shown that gamma-ray




Fig. 2: 2-D reconstruction of points of origin of showers. For small impact parameters, the image has a structure close to that of a circle, whereas with increasing impact parameter it appears more elongated.

on the ground and c) asymmetrical, images are weighted towards their point of origin (Figure 2). The distance between the image centroid and the true point of origin of a shower is represented by the imaging parameter *disp*:

$$disp = 1.85 - 1.85(width/length) \quad (1)$$

from which a unique point of origin can be calculated. As each event represents in itself a unique telescope position in Right Ascension (RA) and Declination (Dec), each point of origin must be translated into an absolute co-ordinate grid. Each point on the grid is then tested for an excess of gamma-ray-like events. The off center properties of the Whipple gamma-ray telescope are well established (Connaughton, et al., 1997b) and are used to determine a flux at any point in the camera.

ACKNOWLEDGEMENTS

We acknowledge the technical support of Emmet Roache. This research is supported by grants from the US Department of Energy and by NASA, by PPARC in the UK, and by Forbairt in Ireland.

REFERENCES

- Akerlof, C.W., et al., 1991, Ap.J. 377, L97
- Cawley, M.F., et al., 1990, Exp. Astron., 1, 185.
- Connaughton, V., et al., 1995, Proc. 24th ICRC, 2, 96.
- Connaughton, V., et al., 1997a, Ap.J., 479, 859.
- Connaughton, V., et al., 1997b, Astroparticle Physics (submitted).
- Hurley, K., Space Science Reviews, v.75, p.43-52.
- Lessard, R.W., et al., 1997, Towards a Major Atmos. Cerenkov Detector - V.
- Reynolds, P.T., et al., 1993, Ap.J. Letters 404, 206